HIV CARE CONTINUUM

Texas DHSHS developed the 2015 HIV/AIDS Care Continuum (HCC) for Dallas EMA, Dallas HSDA and Sherman-Denison HSDA.¹ Along with the Continuum or "cascade," a "Healthier Community" graphical display was developed to further depict linkage and retention in care. The following presents the TDSHS information.

Definitions used in the HCC and Healthier Community calculations:

- HIV+ Individuals at end of 2015 = No. of HIV+ individuals (alive) residing in Texas, Dallas EMA, Dallas HSDA or Sherman-Dennison HSDA at the end of 2015.
- At Least One Visit in 2015 = No. of PLWH with a met need (at least one: medical visit, ART prescription, VL test, or CD-4 test) in 2015, otherwise referred to as "linked to care."
- Retained in Care = No. of PLWH with at least 2 visits or labs, at least 3 months apart or suppressed at end of 2015.
- Achieved Viral Suppression at end of 2015 = No. of PLWH whose last viral load test value of 2015 was <= 200 copies/mL.


In 2015, all three regions reported relatively consistent performance for the four indicators of the HCC.


- Dallas EMA and Dallas HSDA results were nearly identical. Sherman-Dennis reported higher linkage (84%), retention (79%) and suppression rates (68%).
- As a measure of favorable outcomes as a result of HIV care, viral suppression was achieved by 86% to 88% of patients with two or more medical visits in 2015.
- The Dallas EMA, Dallas HSDA and Sherman-Denison HSDA surpassed viral suppression rates when compared to the statewide average.


¹ Sources: Enhanced HIV AIDS Reporting System (eHARS) as of July 2, 2015, Medicaid, ARIES, ADAP, and private payers. Prepared by Program Planning and Evaluation Group, HIV/STD Branch at the Texas Department of State Health Services, August, 2016.

| | PL | WН | At Least (| One Visit | Retaine | d in Care | Suppressed | | % Suppressed of Those Retained |
|---------------------|--------|--------|------------|-----------|------------------|-----------|------------|----------------|---|
| | # | % | # | % | # | % | # | % | % |
| Dallas EMA | 19,793 | 100.0% | 15,664 | 7799%11% | 14, ≩8 ુ% | 72.2% | 12,533 | 633% 3% | 873 %% |
| Dallas HSDA | 19,768 | 100.0% | 15,641 | 79.1% | 14,260 | 72.1% | 12,513 | 63.3% | 87.8% |
| Sherman- Denison | 226 | 100.0% | 189 | 83.6% | 179 | 79.2% | 154 | 68.1% | 86.0% |
| | | | | | | | | | |
| Texas | 82,745 | 100.0% | 63,706 | 77.0% | 57,074 | 69.0% | 48,632 | 59.0% | 85.2% |

HIV Care Continuum Dallas EMA, Dallas HSDA and Sherman-Denison HSDA 2015







<u>Gender</u>

- Across the Dallas EMA, Dallas HSDA, and Sherman-Denison HSDA, similar percentages of male and female PLWH had a met need in 2015.
- 75% of women and 80% of men in the Sherman-Denison HSDA were retained in care, which is higher than 70% of women and 73% of men in both the Dallas EMA and Dallas HSDA.
- Of those retained in care in the Sherman-Denison HSDA, the percentage suppressed was highest for women at 86%; and the Dallas HSDA had the highest for men at 88%. For both genders, the percent suppressed of those retained exceeded 75% for the Dallas EMA, Dallas HSDA, and Sherman-Denison HSDA.

Race/Ethnicity

- Every race/ethnicity category reported over 75% of PLWH a met need in the Dallas EMA, Dallas HSDA, and Sherman-Denison HSDA.
- Sherman-Denison HSDA had the highest percentage of Whites (82%), Blacks (68%) and Hispanics (78%) retained in care. The Dallas EMA had 79% of Whites, 67% of Blacks, and 71% of Hispanics retained in care, which was nearly identical to the percentages in the Dallas HSDA.
- More than half (56%) of Blacks in the Dallas EMA and Dallas HSDA were virally suppressed in 2015, while only 49% of Blacks in Sherman-Denison HSDA were virally suppressed.

<u>Age</u>

- Approximately 80% of PLWH from each age group reported a met need in each geography.
- The percentage suppressed of those retained in care in the Dallas EMA, Dallas HSDA, and Sherman-Denison HSDA exceeded 75% in each age category.

Transmission Mode

- By transmission mode, high percentages of PLWH were identified with met need in 2015.
- In the Dallas EMA, the injection drug-use transmission mode had the lowest percentage of PLWH retained in care (68%). Injection drug-use and MSM/IDU had the lowest number retained in care in the Sherman-Denison HSDA at 72%, while heterosexual transmission was the transmission category in the Dallas HSDA with the lowest percentage retained in care (69%).

HIV Care Continuum by Gender, Race/Ethnicity, Age, Transmission Mode Dallas EMA 2015

| | PLWH | | At Least One Visit | | Retained in Care | | Suppressed | | % Suppressed of those Retained | | |
|---------------------|--------|-------|-----------------------|------|------------------|------|------------|------|--------------------------------------|--|--|
| | # | % | # | % | # | % | # | % | | | |
| ALL PLWH | 19,793 | 100.0 | 15,664 | 79.1 | 14,281 | 72.2 | 12,533 | 63.3 | 80.0% | | |
| Gender | Gender | | | | | | | | | | |
| (F)Female | 3,917 | 19.8 | 3,038 | 77.6 | 2,743 | 70.0 | 2,354 | 60.1 | 77.5% | | |
| (M)Male | 15,876 | 80.2 | 12,626 | 79.5 | 11,538 | 72.7 | 10,179 | 64.1 | 80.6% | | |
| Race/Ethnicity | | | | | | | | | | | |
| White, not Hispanic | 6,245 | 31.6 | 5,205 | 83.3 | 4,904 | 78.5 | 4,492 | 71.9 | 86.3% | | |
| Black, not Hispanic | 8,178 | 41.3 | 6,246 | 76.4 | 5,492 | 67.2 | 4,596 | 56.2 | 73.6% | | |
| Hispanic | 4,372 | 22.1 | 3,369 | 77.1 | 3,112 | 71.2 | 2,773 | 63.4 | 82.3% | | |
| Other | 238 | 1.2 | 180 | 75.6 | 166 | 70.1 | 157 | 66.0 | 87.2% | | |

| | PLWH | | At Least One Visit | | Retained in Care | | Suppressed | | % Suppressed of those Retained |
|----------------------|--------|------|-----------------------|------|------------------|------|------------|------|--------------------------------------|
| | # | % | # | % | # | % | # | % | |
| Unknown | 760 | 3.8 | 664 | 87.4 | 607 | 79.9 | 515 | 67.8 | 77.6% |
| Age | | | | | | | | | |
| 0-24 | 955 | 4.8 | 759 | 80.0 | 581 | 60.8 | 446 | 46.7 | 76.8% |
| 25-34 | 3,848 | 19.4 | 2,995 | 77.8 | 2,563 | 66.6 | 2,093 | 54.4 | 69.9% |
| 35-44 | 4,757 | 24.0 | 3,655 | 76.8 | 3,335 | 70.1 | 2,928 | 61.6 | 80.1% |
| 45-54 | 6,191 | 31.3 | 5,045 | 81.5 | 4,740 | 76.6 | 4,232 | 68.4 | 83.9% |
| 55+ | 4,039 | 20.4 | 3,206 | 79.4 | 3,059 | 75.7 | 2,833 | 70.1 | 88.4% |
| Mode of Transmission | n | | | | | | | | |
| MSM | 13,432 | 67.9 | 10,792 | 80.3 | 9,874 | 73.5 | 8,790 | 65.4 | 81.4% |
| IDU | 1,375 | 6.9 | 1,041 | 75.7 | 939 | 68.3 | 789 | 57.4 | 75.8% |
| MSM/IDU | 809 | 4.1 | 651 | 80.5 | 591 | 73.1 | 472 | 58.3 | 72.5% |
| Heterosexual | 4,024 | 20.3 | 3,062 | 76.1 | 2,770 | 68.8 | 2,400 | 59.6 | 78.4% |
| Pediatric | 131 | 0.7 | 102 | 77.9 | 93 | 71.0 | 69 | 52.7 | 67.6% |
| Adult Other | 22 | 0.1 | 16 | 72.7 | 15 | 68.2 | 12 | 54.5 | 75.0% |

HIV Care Continuum by Gender, Race/Ethnicity, Age, Transmission Mode Dallas HSDA 2015


| | PLWH | | At Least One Visit | | Retained in Care | | Suppressed | | % Suppressed of those Retained |
|--------------------|--------|-------|-----------------------|------|------------------|------|------------|------|--------------------------------------|
| | # | % | # | | # | % | # | % | % |
| All PLWH | 19,768 | 100.0 | 15,641 | 79.1 | 14,260 | 72.1 | 12,513 | 63.3 | 87.7% |
| Gender | | | | | | | | - | |
| (F)Female | 3,928 | 19.9 | 3,049 | 77.6 | 2,753 | 70.1 | 2,362 | 60.1 | 85.8% |
| (M)Male | 15,840 | 80.1 | 12,592 | 79.5 | 11,507 | 72.6 | 10,151 | 64.1 | 88.2% |
| Race/Ethnicity | | | | | | | | | |
| White Non-Hispanic | 6,192 | 31.3 | 5,158 | 83.3 | 4,860 | 78.5 | 4,455 | 71.9 | 91.7% |
| Black Non-Hispanic | 8,202 | 41.5 | 6,268 | 76.4 | 5,512 | 67.2 | 4,610 | 56.2 | 83.6% |
| Hispanic | 4,375 | 22.1 | 3,369 | 77.0 | 3,113 | 71.2 | 2,774 | 63.4 | 89.1% |
| All Other/Unknown | 999 | 5.1 | 846 | 84.7 | 775 | 77.6 | 674 | 67.5 | 87.0% |
| Age | | | | | | | | | |
| < 24 | 956 | 4.8 | 762 | 80.3 | 583 | 61.0 | 448 | 46.9 | 76.8% |
| 25 - 44 | 8,609 | 43.6 | 6,654 | 77.3 | 5,902 | 68.6 | 5,023 | 58.3 | 85.1% |
| > 44 | 10,203 | 51.6 | 8,225 | 80.6 | 7,775 | 76.2 | 7,042 | 69.0 | 90.6% |

| | PLWH | | At Least One Visit | | Retained | in Care | Suppre | ssed | % Suppressed of those Retained | |
|----------------------|----------------------|------|-----------------------|------|----------|---------|--------|------|--------------------------------------|--|
| | # | % | # | | # | % | # | % | % | |
| Mode of Transmission | Mode of Transmission | | | | | | | | | |
| MSM | 13,398 | 67.8 | 10,765 | 80.3 | 9,849 | 73.5 | 8,770 | 65.5 | 89.0% | |
| IDU or MSM/IDU | 2,180 | 11.0 | 1,686 | 77.3 | 1,522 | 69.8 | 1,254 | 57.5 | 82.4% | |
| Heterosexual | 4,038 | 20.4 | 3,072 | 76.1 | 2,781 | 68.9 | 2,407 | 59.6 | 86.6% | |
| All Other | 153 | 0.8 | 118 | 77.1 | 108 | 70.6 | 81 | 52.9 | 75.0% | |

HIV Care Continuum by Gender, Race/Ethnicity, Age, Transmission Mode Sherman-Denison HSDA 2015

| | PLWH | | | At Least One Visit | | Retained in Care | | essed | % Suppressed of those Retained | |
|----------------------|------|-------|-----|-----------------------|-----|---------------------|-----|-------|--------------------------------|--|
| | # | % | # | % | # | % | # | % | % | |
| All PLWH | 226 | 100.0 | 189 | 83.6 | 179 | 79.2 | 154 | 68.1 | 86.0% | |
| Gender | | | | | | | | | | |
| (F)Female | 48 | 21.2 | 37 | 77.1 | 36 | 75.0 | 31 | 64.6 | 86.1% | |
| (M)Male | 178 | 78.8 | 152 | 85.4 | 143 | 81.5 | 123 | 69.1 | 86.0% | |
| Race/Ethnicity | | | | | | | | | | |
| White Non-Hispanic | 146 | 64.6 | 123 | 84.2 | 119 | 81.5 | 103 | 70.5 | 86.6% | |
| Black Non-Hispanic | 41 | 18.1 | 33 | 80.5 | 28 | 68.3 | 20 | 48.8 | 71.4% | |
| Hispanic | 23 | 10.2 | 19 | 82.6 | 18 | 78.3 | 17 | 73.9 | 94.4% | |
| All Other/Unknown | 16 | 7.1 | 14 | 87.5 | 14 | 87.5 | 14 | 87.5 | 100.0% | |
| Age | | | | | | | | | | |
| < 24 | 10 | 4.4 | 10 | 100.0 | 8 | 80.0 | 7 | 70.0 | 87.5% | |
| 25 - 44 | 86 | 38.1 | 70 | 81.4 | 67 | 77.9 | 57 | 66.3 | 85.1% | |
| > 44 | 130 | 57.5 | 109 | 83.8 | 104 | 80.0 | 90 | 69.2 | 86.5% | |
| Mode of Transmission | | | | | | | | | | |
| MSM | 117 | 51.8 | 103 | 88.0 | 97 | 82.9 | 84 | 71.8 | 86.6% | |
| IDU or MSM/IDU | 54 | 23.9 | 41 | 75.9 | 39 | 72.2 | 31 | 57.4 | 79.5% | |
| Heterosexual | 51 | 22.6 | 43 | 84.3 | 41 | 80.4 | 37 | 72.5 | 90.2% | |
| All Other | 4 | 2.0 | 2 | 50.0 | 2 | 50.0 | 2 | 50.0 | 100.0% | |

2015 Care Continuum





HEALTHIER COMMUNITY



79% of PLWH had at least one episode of HIV care & treatment. This means roughly 8 out of 10 PLWH were in care.



72% of PLWH were retained in care (2 episodes of HIV care & treatment across the year). This means that roughly 7 out of 10 PLWH were retained in care

İİİİİİİİİİİ


63% of PLWH were virally suppressed. This means that roughly 6 out of 10 PLWH were virally suppressed




Of those 7 out of 10 PLWH who were retained in care, 88%, or roughly 6 of those 7 PLWH, were virally suppressed.

2015 Continuum of Care, Parity Table

| | | ŴН | At leas vis | it | Retaine Care | 2 | % retained if any care | Suppressed | | % suppressed of those retained |
|---------------|--------|------|----------------|-------------|-----------------|------------------|---------------------------|------------|-----|--------------------------------------|
| | # | % | # | % | # | % | % | # | % | % |
| All PLWH | 19,768 | 100% | 15,641 | 79 % | 14,260 | <mark>72%</mark> | 91% | 12,513 | 635 | <mark>88%</mark> |
| Female | 3,928 | 20% | 3,049 | 78% | 2,753 | <mark>70%</mark> | 90% | 2,362 | 60% | <mark>86%</mark> |
| Male | 15,840 | 80% | 12,592 | 79% | 11,507 | <mark>73%</mark> | 91% | 10,151 | 64% | <mark>885</mark> |
| White | 6,192 | 31% | 5,158 | 83% | 4,860 | <mark>78%</mark> | 94% | 4,455 | 72% | <mark>92%</mark> |
| Black | 8,202 | 41% | 6,268 | 76% | 5,512 | <mark>67%</mark> | 88% | 4,610 | 56% | <mark>84%</mark> |
| Latinx | 4,375 | 22% | 3,369 | 77% | 3,113 | <mark>71%</mark> | 92% | 2,774 | 63% | <mark>89%</mark> |
| <=24 | 956 | 5% | 762 | 80% | 583 | <mark>61%</mark> | 77% | 448 | 47% | <mark>77%</mark> |
| 25 – 44 | 8,609 | 44% | 6,654 | 77% | 5,902 | <mark>69%</mark> | 89% | 5,023 | 58% | <mark>85%</mark> |
| >= 45 | 10,203 | 52% | 8,225 | 81% | 7,775 | <mark>76%</mark> | 95% | 7,042 | 69% | <mark>91%</mark> |
| MSM | 13,398 | 68% | 10,765 | 80% | 9,849 | <mark>74%</mark> | 91% | 8,770 | 65% | <mark>89%</mark> |
| IDU | 2,180 | 11% | 1,686 | 77% | 1,522 | <mark>70%</mark> | 90% | 1,254 | 58% | <mark>82%</mark> |
| Heterosexual | 4,038 | 20% | 3,072 | 76% | 2,781 | <mark>69%</mark> | 91% | 2,407 | 60% | <mark>87%</mark> |
| White MSM | 5,185 | 26% | 4,360 | 84% | 4,115 | <mark>79%</mark> | 94% | 3,807 | 73% | <mark>93%</mark> |
| Black MSM | 4,271 | 22% | 3,288 | 77% | 2,854 | <mark>67%</mark> | 87% | 2,390 | 56% | <mark>84%</mark> |
| Latino MSM | 3,296 | 17% | 2,570 | 78% | 2,376 | <mark>72%</mark> | 92% | 2,132 | 65% | <mark>90%</mark> |
| Black Females | 2,648 | 13% | 2,027 | 77% | 1,810 | <mark>68%</mark> | 89% | 1,543 | 58% | <mark>85%</mark> |



2015 Continuum of Care, Parity Bar Charts



Targets

85% PLWH retained in HIV care & treatment

81% of those retained are virally suppressed

| | PLV | | Retained Care | | 85% retained goal | Gap | Suppressed | 81% suppressed goal | Gap |
|------------------|--------|------|------------------|-----|-------------------------|-------|------------|---------------------------|-------|
| | # | % | # | % | # | # | # | # | # |
| All PLWH | 19,768 | 100% | 14,260 | 72% | 16,803 | 2,543 | 12,513 | 13,610 | 1,097 |
| Female | 3,928 | 20% | 2,753 | 70% | 3,339 | 586 | 2,362 | 2,705 | 343 |
| Male | 15,840 | 80% | 11,507 | 73% | 13,464 | 1,957 | 10,151 | 10,906 | 755 |
| White | 6,192 | 31% | 4,860 | 78% | 5,263 | 403 | 4,455 | 4,263 | -192 |
| Black | 8,202 | 41% | 5,512 | 67% | 6,972 | 1,460 | 4,610 | 5,647 | 1,037 |
| Latinx | 4,375 | 22% | 3,113 | 71% | 3,719 | 606 | 2,774 | 3,012 | 238 |
| <=24 | 956 | 5% | 583 | 61% | 813 | 230 | 448 | 659 | 211 |
| 25 - 44 | 8,609 | 44% | 5,902 | 69% | 7,318 | 1,416 | 5,023 | 5,928 | 905 |
| >= 45 | 10,203 | 52% | 7,775 | 76% | 8,673 | 898 | 7,042 | 7,025 | -17 |
| MSM | 13,398 | 68% | 9,849 | 74% | 11,388 | 1,539 | 8,770 | 9,224 | 454 |
| IDU | 2,180 | 11% | 1,522 | 70% | 1,853 | 331 | 1,254 | 1,501 | 247 |
| Heterosexual | 4,038 | 20% | 2,781 | 69% | 3,432 | 651 | 2,407 | 2,780 | 373 |
| White MSM | 5,185 | 26% | 4,115 | 79% | 4,407 | 292 | 3,807 | 3,570 | -237 |
| Black MSM | 4,271 | 22% | 2,854 | 67% | 3,630 | 776 | 2,390 | 2,940 | 550 |
| Latino MSM | 3,296 | 17% | 2,376 | 72% | 2,801 | 425 | 2,132 | 2,269 | 137 |
| Black Females | 2,648 | 13% | 1,810 | 68% | 2,251 | 441 | 1,543 | 1,823 | 280 |